

Acknowledgments

We thank N. Bornfeld, B. Lorenz, U. Oehlschläger, and S. Schüler, for referral of patients, and E. Passarge for continuous support.

MARTINA KLUTZ, BERNHARD HORSTHEMKE, AND
DIETMAR R. LOHMANN

*Institut für Humangenetik,
Universitätsklinikum Essen,
Essen, Germany*

Electronic-Database Information

URL for data in this article is as follows:

RB1 gene mutation database, <http://home.kamp.net/home/dr.lohmann>

References

- Bremner R, Du DC, Connolly-Wilson MJ, Bridge P, Ahmad KF, Mostachfi H, Rushlow D, et al (1997) Deletion of RB exons 24 and 25 causes low-penetrance retinoblastoma. *Am J Hum Genet* 61:556–70
- Draper GJ, Sanders BM, Brownbill PA, Hawkins MM (1992) Patterns of risk of hereditary retinoblastoma and applications to genetic counselling. *Br J Cancer* 66:211–219
- Gallie BL (1997) Predictive testing for retinoblastoma comes of age. *Am J Hum Genet* 61:279–281
- Greger V, Debus N, Lohmann D, Höpping W, Passarge E, Horsthemke B (1994) Frequency and parental origin of hypermethylated RB1 alleles in retinoblastoma. *Hum Genet* 94:491–496
- Hu QJ, Dyson N, Harlow E (1990) The regions of the retinoblastoma protein needed for binding to adenovirus E1A or SV40 large T antigen are common sites for mutations. *EMBO J* 9:1147–1155
- Lohmann DR, Brandt B, Höpping W, Passarge E, Horsthemke B (1996) Spectrum of RB1 germ-line mutations in hereditary retinoblastoma. *Am J Hum Genet* 58:940–949
- Lohmann DR, Gerick M, Brandt B, Oehlschläger U, Lorenz B, Passarge E, Horsthemke B (1997) Constitutional RB1-gene mutations in patients with isolated unilateral retinoblastoma. *Am J Hum Genet* 61:282–294
- Otterson GA, Chen WD, Coxon AB, Khleif SN, Kaye FJ (1997) Incomplete penetrance of familial retinoblastoma linked to germ-line mutations that result in partial loss of RB function. *Proc Natl Acad Sci USA* 94:12036–12040
- Sippel KC, Fraioli RE, Smith GD, Schalkoff ME, Sutherland J, Gallie BL, Dryja TP (1998) Frequency of somatic and germ-line mosaicism in retinoblastoma: implications for genetic counseling. *Am J Hum Genet* 62:610–619
- Toguchida J, McGee TL, Paterson JC, Eagle JR, Tucker S, Yandell DW, Dryja TP (1993) Complete genomic sequence of the human retinoblastoma susceptibility gene. *Genomics* 17:535–543
- Vogel F (1979) Genetics of retinoblastoma. *Hum Genet* 52:1–54

Yandell DW, Campbell TA, Dayton SH, Petersen R, Walton D, Little JB, McConkie-Rosell A, et al (1989) Oncogenic point mutations in the human retinoblastoma gene: their application to genetic counseling. *N Engl J Med* 321:1689–1695

Address for correspondence and reprints: Dr. Dietmar R. Lohmann, Institut für Humangenetik, Hufelandstrasse 55, D-45122 Essen, Germany. E-mail: dr.lohmann@uni-essen.de

Am. J. Hum. Genet. 64:668–669, 1999

TDT Clarification

To the Editor:

A potentially misleading statement occurs in the invited editorial “The TDT and other family-based tests for linkage disequilibrium and association” by Spielman and Ewens (59:983–989), published in the November 1996 issue of the *Journal*. We wish to make the following clarification.

In discussing some issues that arise when the transmission/disequilibrium test (TDT) is used in families where genotype data are unavailable for one parent, we noted the finding of Curtis and Sham (1995) that, when there is a single affected offspring who is homozygous for an allele present in the available (heterozygous) parent, the TDT gives a biased result and should not be used. We then stated that “[w]hen there is more than one offspring in the sibship, it sometimes will be possible to deduce that the unavailable parent [is also heterozygous], and, in these cases, we may proceed as though this [reconstructed] genotype were known directly” (Spielman and Ewens 1996, p. 987).

We should have emphasized that this claim assumes that the reconstruction is done from the genotypes of unaffected offspring. A bias will usually arise in the TDT statistic if the reconstruction uses, in whole or in part, genotype data from affected offspring whose genotypes are then used in the TDT. A bias can also arise when both parental genotypes are reconstructed from the genotypes of the offspring.

The bias resulting from reconstruction occurs for a reason different from that noted by Curtis and Sham (1995). Since reconstruction is possible only with certain offspring genotype combinations, families in which parental genotypes can be reconstructed are not random as far as offspring genotypes are concerned. This restriction leads to a bias if the offspring genotype data that were used for reconstruction are also used in the TDT.

Knapp (in press) has calculated the size of this bias in a number of important cases and has thus been able to establish more general TDT tests where this bias is allowed for.

Acknowledgments

We thank Michael Knapp for many valuable discussions concerning reconstruction of parental genotypes.

RICHARD S. SPIELMAN¹ AND WARREN J. EWENS²
Departments of ¹Genetics and ²Biology, University of Pennsylvania, Philadelphia

References

- Curtis D, Sham PC (1995) A note on the application of the transmission disequilibrium test when a parent is missing. *Am J Hum Genet* 56:811–812
- Knapp M. The transmission/disequilibrium test (TDT) and parental genotype reconstruction: the reconstruction combined TDT (RC-TDT). *Am J Hum Genet* (in press)
- Spielman RS, Ewens WJ (1996) The TDT and other family-based tests for linkage disequilibrium and association. *Am J Hum Genet* 59:983–989

Address for correspondence and reprints: Dr. Warren J. Ewens, Leidy Laboratory of Biology, Department of Biology, University of Pennsylvania, Philadelphia, PA 19104-6018. E-mail: wewens@sas.upenn.edu